

PARALLELOPIPEDS OF POSITIVE RANK TWISTS OF ELLIPTIC CURVES

BO-HAE IM AND MICHAEL LARSEN

ABSTRACT. For every $n \in \mathbb{N}$ there exists an elliptic curve $E/\mathbb{Q}$ and an n -dimensional subspace V of $\mathbb{Q}^\times/(\mathbb{Q}^\times)^2$ such that for all $v \in V$, the quadratic twist E_v has positive rank.

1. INTRODUCTION

Let E be an elliptic curve over a number field K . For every element $v \in K^\times/(K^\times)^2$, there is a well-defined quadratic twist of E associated to v , which we denote E_v . The main result of this paper is the following:

Theorem 1. *For every $n \in \mathbb{N}$ there exists an elliptic curve $E/\mathbb{Q}$ and an n -dimensional subspace V of $\mathbb{Q}^\times/(\mathbb{Q}^\times)^2$ such that for all $v \in V$, the quadratic twist E_v has positive rank.*

We can express this theorem slightly differently, along the lines of [IL]. If L/K is a finite Galois extension of number fields, with group G and E/K is an elliptic curve, then G acts on $V := E(L) \otimes \mathbb{Q}$. If W is any G -subrepresentation of V , we say that the pair (G, W) is *Mordell-Weil over K* . Theorem 1 now asserts that the regular representation of every elementary 2-group is Mordell-Weil over $\mathbb{Q}$. We remark that T. Dokchitser and V. Dokchitser [DD] have given examples of number fields K such that, modulo the Birch-Swinnerton-Dyer conjecture, every regular representation of an elementary 2-group is Mordell-Weil over K in dimension 1. In fact, something stronger is true: their examples have the remarkable property that (under Birch-Swinnerton-Dyer) *every* quadratic twist has positive rank. (Of course, K cannot be $\mathbb{Q}$.)

Our proof depends on Vatsal's theorem [Va] which asserts that for the elliptic curve $X_0(19)$ the set of positive integers d such that E_d has rank 1 has positive density.

2000 *Mathematics Subject Classification.* 11G05.

Michael Larsen was partially supported by NSF grant DMS-0800705.

2. SOME MULTIPLICATIVE COMBINATORICS

Definition 2. Let n be a positive integer. We say that a subset Π of $\mathbb{N}$ is a strict n -parallelopiped if we have

$$\Pi = \{c \prod_{i \in I} a_i \mid I \subset \{1, 2, \dots, n\}\}$$

for some rational numbers $a_i, c \in \mathbb{Q}^*$ such that the images of $a_1, \dots, a_n$ are linearly independent in the $\mathbb{F}_2$ -vector space $\mathbb{Q}^*/\mathbb{Q}^{*2}$.

The main result of this section is the following:

Theorem 3. Any set S of positive integers of positive lower density contains a strict n -parallelopiped for each positive integer n .

For any subset S of $\mathbb{N}$, we define the function

$$f_S(t) := \sum_{n \in S} e^{-nt}$$

on the domain $(0, \infty)$. Thus, $S \subset T$ implies that for all t , $f_S(t) \leq f_T(t)$. In particular,

$$(1) \quad f_S(t) \leq f_{\mathbb{N}}(t) = \frac{1}{1 - e^{-t}} \leq 2 \max(1/t, 1).$$

More generally, this principle applies to multisets. If $S_1, \dots, S_n \subset \mathbb{N}$, then for all $t \in (0, \infty)$, the first two Bonferroni inequalities imply

$$(2) \quad f_{S_1 \cup \dots \cup S_n}(t) \leq \sum_{1 \leq i \leq n} f_{S_i}(t)$$

and

$$(3) \quad \sum_{1 \leq i \leq n} f_{S_i}(t) - \sum_{1 \leq i < j \leq n} f_{S_i \cap S_j}(t) \leq f_{S_1 \cup \dots \cup S_n}(t).$$

If $a \leq b$, we define

$$P_{a,b} = \prod_{a \leq p \leq b} p,$$

where the product is taken over primes. For any positive integer m , we define

$$R(m) := \{n \in \mathbb{N} \mid (m, n) = 1\}$$

the set of positive integers relatively prime to m .

Lemma 4. For any $S \subset \mathbb{N}$ and $b \geq a > 1$ and for all $t \in (0, \infty)$, we have

$$f_{S \cap R(P_{a,b})} = \sum_{\{n \in S \mid (n, P_{a,b}) = 1\}} e^{-nt} \leq t^{-1} \prod_{a \leq p \leq b} (1 - p^{-1}).$$

Proof. Clearly

$$f_{S \cap R(P_{a,b})}(t) \leq f_{R_{a,b}}(t) = \sum_{i=0}^{\infty} \sum_{\{n \in (iP_{a,b}, (i+1)P_{a,b}) \mid (n, P_{a,b})=1\}} e^{-nt}.$$

As

$$\sum_{\{n \in (iP_{a,b}, (i+1)P_{a,b}) \mid (n, P_{a,b})=1\}} e^{-nt} \leq \phi(P_{a,b}) e^{-iP_{a,b}t},$$

we have

$$f_{R_{a,b}}(t) \leq \frac{\phi(P_{a,b})}{1 - e^{-P_{a,b}t}} \leq \frac{\phi(P_{a,b})}{tP_{a,b}} = t^{-1} \prod_{a \leq p \leq b} (1 - p^{-1}).$$

□

For any $S \subset \mathbb{N}$ we define

$$D(S) := \limsup_{T \rightarrow \infty} \frac{1}{\log T} \int_{1/T}^1 f_S(t) dt.$$

Lemma 5. *For all subsets S of $\mathbb{N}$, let $d(S)$ denote the lower density of S . Then*

$$D(S) \geq d(S).$$

Proof. For all $\epsilon > 0$, there exists N such that for all $n \geq N$,

$$|S \cap [1, n]| > (d(S) - \epsilon)n.$$

Therefore, for all $t > 0$,

$$f_S(t) > \sum_{n=N}^{\infty} (d(S) - \epsilon) e^{-nt} = \frac{e^{-Nt}(d(S) - \epsilon)}{1 - e^{-t}} \geq \frac{e^{-Nt}(d(S) - \epsilon)}{t}.$$

Thus, for $T > N/\epsilon$, we have

$$\begin{aligned} \frac{1}{\log T} \int_{1/T}^1 f_S(t) dt &\geq \frac{1}{\log T} \int_{1/T}^{\epsilon/N} \frac{e^{-\epsilon}(d(S) - \epsilon)}{t} dt \\ &\geq \left(1 - \frac{\log N/\epsilon}{\log T}\right) e^{-\epsilon}(d(S) - \epsilon). \end{aligned}$$

Taking $\limsup_{T \rightarrow \infty}$, we conclude that for all $\epsilon > 0$,

$$D(S) \geq e^{-\epsilon}(d(S) - \epsilon),$$

which implies the lemma. □

For any subset S of $\mathbb{N}$ and any positive integer m , let

$$S_m := \mathbb{N} \cap m^{-1}S = \{n \in \mathbb{N} \mid mn \in S.\}$$

Proposition 6. *If S is a subset of $\mathbb{N}$ with $D(S) > 0$, and N is a positive integer, there exist primes $q > p > N$ such that*

$$D(S_p \cap S_q) > 0.$$

Proof. Let a be an integer larger than $\max(N, \frac{12}{D(S)})$. As

$$\prod_{p \geq a \text{ prime}} (1 - p^{-1}) = \prod_{p < a \text{ prime}} (1 - p^{-1})^{-1} \prod_{p \text{ prime}} (1 - p^{-1}) = 0,$$

we may choose b such that

$$\prod_{\substack{a \leq p \leq b \\ p \text{ prime}}} (1 - p^{-1}) \leq \frac{D(S)}{4}.$$

Writing

$$S = (S \cap R(P_{a,b})) \cup \bigcup_{a \leq p \leq b} pS_p,$$

and applying (2) and Lemma 4, we obtain

$$\begin{aligned} \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} f_{S_p}(pt) &= \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} f_{pS_p}(t) \geq f_S(t) - f_{S \cap R(P_{a,b})}(t) \\ (4) \qquad \qquad \qquad &\geq f_S(t) - t^{-1} \prod_{\substack{a \leq p \leq b \\ p \text{ prime}}} (1 - p^{-1}). \end{aligned}$$

Let X denote the set of values T such that

$$\frac{1}{\log T} \int_{1/T}^1 f_S(t) dt \geq \frac{3D(S)}{4}.$$

By definition of $D(S)$, X is unbounded. Choosing $T \in X$ and integrating (4), we obtain

$$\begin{aligned} (5) \qquad \frac{1}{\log T} \int_{1/T}^1 \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} f_{S_p}(pt) dt &\geq \frac{1}{\log T} \int_{1/T}^1 f_S(t) dt - \prod_{\substack{a \leq p \leq b \\ p \text{ prime}}} (1 - p^{-1}) \\ &\geq \frac{D(S)}{2}. \end{aligned}$$

Interchanging integral and sum and substituting $u = pt$, we obtain

$$\frac{1}{\log T} \int_{1/T}^1 \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} f_{S_p}(pt) dt = \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \frac{1}{p \log T} \int_{p/T}^p f_{S_p}(u) du.$$

By (1), this can expressed as

$$\sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \frac{1}{p \log T} \int_{1/T}^1 f_{S_p}(u) du + O\left(\frac{1}{\log T}\right).$$

Combining this with (5) and choosing T sufficiently large, we obtain

$$\frac{1}{\log T} \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \int_{1/T}^1 f_{S_p}(u) du \geq \frac{1}{\log T} \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \frac{a}{p} \int_{1/T}^1 f_{S_p}(u) du \geq \frac{aD(S)}{3}.$$

By definition of a , this means

$$(6) \quad \frac{1}{\log T} \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \int_{1/T}^1 f_{S_p}(u) du \geq 4.$$

Let

$$S' = \bigcup_{a \leq p \leq b} S_p.$$

We integrate the inequality (3) from $1/T$ to 1 and interchange sum and integral to obtain

$$(7) \quad \sum_{\substack{a \leq p < q \leq b \\ p, q \text{ prime}}} \frac{1}{\log T} \int_{1/T}^1 f_{S_p \cap S_q}(u) du \geq \frac{1}{\log T} \sum_{\substack{a \leq p \leq b \\ p \text{ prime}}} \int_{1/T}^1 f_{S_p}(u) du - \frac{1}{\log T} \int_{1/T}^1 f_{S'}(u) du.$$

Applying (1) to S' and integrating, we get

$$\frac{1}{\log T} \int_{1/T}^1 f_{S'}(u) du \leq 2.$$

Combining this with (6) and (7), we get

$$\sum_{\substack{a \leq p < q \leq b \\ p, q \text{ prime}}} \frac{1}{\log T} \int_{1/T}^1 f_{S_p \cap S_q}(u) du \geq 2,$$

which implies

$$D(S_p \cap S_q) \geq \frac{1}{(b-a)(1+b-a)}.$$

for some p and q . □

The following theorem is a strengthened version of Theorem 3.

Theorem 7. *Let Σ denote a finite set of primes and S a subset of $\mathbb{N}$ such that $D(S) > 0$. Then there exist a rational number c and an infinite set of pairwise distinct vectors $(a_{j,1}, \dots, a_{j,n}) \in \mathbb{Q}^n$ such that for each j ,*

$$\{c \prod_{i \in I} a_{j,i} \mid I \subset \{1, 2, \dots, n\}\} \subset S,$$

the numerators and denominators of the $a_{j,i}$ are not divisible by any prime in Σ , and the images of $a_{j,1}, \dots, a_{j,n}$ are linearly independent in $\mathbb{Q}^/\mathbb{Q}^{*2}$.*

Proof. We use induction on n . The base case $n = 1$ is trivial. Suppose the theorem is true for some $n \geq 1$. Let N be the maximum of all primes in Σ . By Proposition 6, there exist primes $q > p > N$ such that $D(S_p \cap S_q) > 0$. By the induction hypothesis for $S_p \cap S_q$ and $\Sigma' := \Sigma \cup \{p, q\}$, there exist c and $a_{j,i}$ prime to Σ' such that the $a_{j,1}, \dots, a_{j,n}$ are independent in $\mathbb{Q}^*/\mathbb{Q}^{*2}$ and for all j ,

$$\{c \prod_{i \in I} a_{j,i} \mid I \subset \{1, 2, \dots, n\}\} \subset S_p \cap S_q.$$

Setting $c' := pc$, $a'_{j,1} := a_{j,1}, \dots, a'_{j,n} := a_{j,n}$, and $a'_{j,n+1} := q/p$, we have

$$\{c' \prod_{i \in I} a'_{j,i} \mid I \subset \{1, 2, \dots, n+1\}\} \subset S,$$

and the $a'_{j,i}$ are prime to Σ . □

3. THE MAIN THEOREM

We prove Theorem 1.

Proof of Theorem 1. Let E be the elliptic curve $X_0(19)$ and

$$S := \{d \in \mathbb{N} \mid \text{rank}(E_d) > 0\}.$$

Then, Vatsal's theorem [Va] implies that S has positive lower density.

Let $n \in \mathbb{N}$ be given. By Theorem 3, there exist $c, a_1, \dots, a_n \in \mathbb{Q}^*$ such that the images of a_i are linearly independent in $\mathbb{Q}^*/\mathbb{Q}^{*2}$ and such that

$$\{c \prod_{i \in I} a_i \mid I \subset \{1, 2, \dots, n\}\} \subset S.$$

If we let V be a subspace of $\mathbb{Q}^*/\mathbb{Q}^{*2}$ generated by $a_1, \dots, a_n$, then it is of dimension n and for each $v \in V$, the rank of the twist of E_c by v (which is E_{cv}) is positive.

REFERENCES

- [DD] Dokchitser, Tim; Dokchitser, Vladimir: Elliptic curves with all quadratic twists of positive rank. *Acta Arith.* **137** (2009), no. 2, 193–197.
- [IL] Im, Bo-Hae; Larsen, Michael: Realizing rational representations in Mordell-Weil groups, preprint.
- [Va] Vatsal, V.: Rank-one twists of a certain elliptic curve. *Math. Ann.* **311** (1998), no. 4, 791–794.

DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, 221, HEUKSEOK-DONG, DONGJAK-GU, SEOUL, 156-756, SOUTH KOREA
E-mail address: bohaeim@gmail.com

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405, USA
E-mail address: larsen@math.indiana.edu